

Andri

Page 1

From: "Shami Goats" <info@shamigoats.com.cy>
To: "Asreda Corporation" <asreda.csb@gmail.com>
Sent: Τρίτη, 15 Σεπτεμβρίου 2009 1:11 μμ
Subject: Re: Re:

Dear Mr Shariff,

Let me know what to do with the cages. We must hurry as we have to make the shipment in October.
Regarding the food. Is not need to eat or drink during the flight. They are going to have a lot of stress so is better not to eat or drink.
They will eat and drink before flight.
At arrival you will give them straw and water and slowly slowly you will give them only straw and barley. For the first 24 hours you will give them hay and mixture of barley and corn maximum half kilo per goat.
The first week you will give them barley and corn to one kilo. After this week you will deduct 100 gr of this food and replace it with the Malaysia food every day. In a week you will give them half barley with corn and half Malaysia food. In approximately 3 weeks you will give them only Malaysia food.

Best Regards
Andri

For Neoclis Antoniadis.

Y.N. ANTONIADES & SONS LTD

P.O.BOX 51611
3507 LIMASSOL
CYPRUS

TEL.: 00357 25 392222
FAX: 00357 25 390885

PROPORTION FOR 1 TONE (1000KG) GOATS

	PREGNANT	MILKY	FUTURE PRODUCTIVE FEMALES
BARLEY	500KG	470KG	750KG
CORN	100KG	200KG	95KG
BRAN	50KG	100KG	20KG
CAROB	100KG		
SOYA BEAN	200KG	180KG	110KG
VITAMINS	2KG	2KG	1KG
CALSIUM	20KG	20KG	10KG
FOSFORO DIASV.	18KG	18KG	9KG
SALT	10KG	10KG	5KG
TOTAL	1000KG	1000KG	1000KG

For Each animals:

a) Milking Animals:

2 - 2 ½ kilogram of animal food for goats is enough for 4 ltrs of milk, for animals that they produce more than 4 ltrs of milk for every 1 ltrs of milk extra ½ kilogram of food.

b) Pregnant 2 - 3 months 600 grams

Pregnant 3 - 4 months 800 grams

To the 5th month of pregnancy 1Kilo

c) For future productive female 750 grams up to 1 kilogram.

Y.N.ANTONIADES AND SONS LTD

P.O.BOX 51611

TEL.25 392222

FAX: 25-390885

LIMASSOL-CYPRUS

Basic Vaccinations Program for Goats

6 weeks before parturition

Dialuene 2 ml/sc (Clostridium Perfigens type B,C,D Clostridium Serticum,
Tetani, Clostridium Novyi, clostridium shauvue).
E Coli(K99)

3 weeks before parturition

Dialuene 2 ml/sc
E Coli

Before matings 20 days

Clamydia Abortions:
Vit E+SE

Kids

Vit E+SE at the 2 days
Dialuend P 2ml/sc at 20-30 days
Baycox per OS at 20-30 days

Dialuene P 2ml/sc at 50-60 days
Oramec 5ml per OS at 50-60 days

Y.N.ANTONIADES AND SONS LTD

P.O.BOX 51611

TEL.25 392222

FAX: 25-390885

LIMASSOL-CYPRUS

Vaccination and Defence program for Goats and Kids

Adult Goats

Enterotoxaemia type D: Every Spring and Autumn

Chlamydia Abortion: One month before mating

Colibacillosis: 6 weeks and 3 weeks before parturition

Vit.E + Selenium: 3 weeks before parturition

Young Kids

Vit.E + Selenium: 2nd or 3rd day of life

Enterotoxaemia taupe D:

If the Kids come from vaccinated mothers you vaccinate of the age of one Month and you repeat vaccination 3 weeks later.

If the kids come from unvaccinated mothers vaccination must be done of The age of 2 weeks and you repeat 3 weeks later.

Coccidiostatic: The feed given to the kids should contain coccidiostatic (Avatec e.t.c.) till the age of 6 months .

In case Kids will be slaughtered stop coccidiostatic one month before.

Additional comment: In case there are specific health problem in the flock f.e

Pneumonia or paratuberculosis etc then additional vaccination against These diseases are needed.